

Leading-twist pion and kaon distribution amplitudes in the gauge-invariant nonlocal chiral quark model from the instanton vacuum

Seung-il Nam^{*} and Hyun-Chul Kim[†]

*Department of Physics and Nuclear Physics & Radiation Technology Institute (NuRI),
Pusan National University, Busan 609-735, Republic of Korea*

(Dated: September, 2006)

Abstract

We investigate the leading-twist light-cone distribution amplitudes for the pion and kaon based on the *gauge-invariant* nonlocal chiral quark model from the instanton vacuum in the presence of external axial-vector currents. We find that the nonlocal contribution from the gauge invariance has much effects on the pion distribution amplitudes, while it changes mildly the kaon ones. We also study the Gegenbauer moments of the distribution amplitudes and compare them with the empirical analysis of the CLEO data.

PACS numbers: 11.15.Tk, 14.40.Aq

Keywords: Meson distribution amplitudes, instanton vacuum, gauge invariance of the nonlocal interaction

^{*}Electronic address: sinam@pusan.ac.kr

[†]Electronic address: hchkim@pusan.ac.kr

I. INTRODUCTION

The meson light-cone distribution amplitude (DA) provides essential information on the nonperturbative structure of mesons. In particular, the leading-twist meson DAs play a role of input for describing hard exclusive reactions due to factorization theorems [1, 2, 3, 4]. In particular, the pion DA has been investigated extensively in various theoretical approaches: For example, in the QCD sum rules (QCDSR) [4, 5, 6, 7, 8, 9], in lattice QCD (LQCD) [10, 11], in the chiral quark model (χ QM) from the instanton vacuum [12, 13, 14], in the NJL models [15, 16, 17] and so on. Recently, the CLEO experiment [18] has measured the $\gamma^*\gamma \rightarrow \pi$ transition form factor, which gives criteria for judging the existing theoretical pion DAs [19]. Bakulev *et al.* [20, 21] have carried out similar analyses.

In the previous work [14], we have investigated the leading-twist pion and kaon DAs within the framework of the nonlocal χ QM from the instanton vacuum with SU(3) symmetry breaking explicitly taken into account. However, it is well known that the nonlocal interaction breaks the conservation of the vector and axial-vector currents [22, 23, 24]. The nonlocal χ QM from the instanton vacuum suffers from the same difficulty due to the zero-mode approximation [24]. Since the leading-twist meson DAs involve the axial-vector operator, one has to deal with the problem of the current conservation. While Ref. [24] proposed a systematic way to remedy this problem of the current conservation, one has to handle the integral equation. Refs. [25, 26] derived the light-quark partition function in the presence of the external gauge fields. With this gauged partition function, it was shown that the low-energy theorem for the transition from two-photon state to the vacuum via the axial anomaly was satisfied [25]. Moreover, the magnetic susceptibility of the QCD vacuum was properly obtained [26].

The conservation of the axial-vector current concerning the leading-twist pion DA was already discussed in Refs. [17, 27] by using the instanton-motivated separable nonlocal kernel for the effective action and the NJL model with the Pauli-Villars regularization, respectively. It was pointed out that the nonlocal contributions are about 30% of the local one [27]. In the present work, we want to investigate the leading-twist pion and kaon DAs, employing the method developed in Refs. [25, 26] with the conservation of the axial-vector current considered. We will show that the nonlocal part of the pion DAs arising from the current conservation plays an important role in improving the results of the corresponding Gegenbauer moments.

The present work is organized as follows: In Section II we show how to make the nonlocal effective chiral action gauge-invariant from the instanton vacuum. In Section III we show how to calculate the leading-twist pion and kaon distribution amplitudes in the present approach. In Section IV we discuss numerical results. In Section V we summarize the present work and draw conclusions.

II. GAUGE-INVARIANT EFFECTIVE CHIRAL ACTION

In this Section, we briefly review how one can make the low-energy effective QCD partition function from the instanton vacuum in the presence of external axial-vector fields, following the approach developed in Ref. [25, 26]. The partition function derived from the instanton vacuum [28, 29] can be written in Euclidean space:

$$\mathcal{Z} = \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{D}\mathcal{M} \exp \int d^4x \left[\bar{\psi}_f^\dagger(x) (i\not{\partial} + im_f) \psi_f(x) \right]$$

$$+i \int \frac{d^4k d^4p}{(2\pi)^8} e^{-i(k-p)\cdot x} \psi_f^\dagger(k) \sqrt{M_f(k)} U_{fg}^{\gamma_5} \sqrt{M_g(p)} \psi_g(p) \Big], \quad (1)$$

where ψ_f , m_f , and M_f denote the quark fields, the current quark mass, and the momentum-dependent dynamical quark mass with flavor f , respectively. The U^{γ_5} or $\mathcal{M}$ represents the pseudo-Goldstone field defined as follows:

$$U^{\gamma_5} = U(x) \frac{1 + \gamma_5}{2} + U^\dagger(x) \frac{1 - \gamma_5}{2} = 1 + \frac{i}{F_{\mathcal{M}}} \gamma_5 \mathcal{M}^a \lambda^a - \frac{1}{2F_{\mathcal{M}}^2} \mathcal{M}^2 \dots, \quad (2)$$

where λ stands for the 3×3 Gell-Mann matrix. Musakhanov [30, 31] improved the partition function in Eq. (1) by taking into account effects of flavor SU(3) symmetry breaking, so that the dynamical quark mass acquires the contribution of the m_f corrections:

$$M_f(k) = M_0 F^2(k) \left[\sqrt{1 + \frac{m_f^2}{d^2}} - \frac{m_f}{d} \right], \quad (3)$$

where M_0 is the dynamical quark mass with zero momentum transfer in the chiral limit. Its value is determined by the saddle-point equation: $M_0 \simeq 350$ MeV. $F(k)$ is the momentum-dependent part which arises from the Fourier transform of the fermionic zero-mode solutions in the instantons. Note that we parameterize $M_0(k)$ as a simple-pole type to make the analytic continuation to Minkowski space easy.

The momentum-dependent dynamical quark mass $M_f(p)$ breaks the conservation of the Nöther currents, i.e. it violates Ward-Takahashi identities. We want to show how to remedy this problem in the following. We first define the total quark propagator $\tilde{S}$ in the presence of the instanton ensemble A and external axial-vector field $a_\mu = a_\mu^a \lambda^a / 2$ and the quark propagator $\tilde{S}_i$ with a single instanton A_i as well as the a_μ :

$$\tilde{S} = \frac{1}{i\not{\partial} + g\not{A} + \not{\phi}\gamma_5 + i\hat{m}}, \quad \tilde{S}_i = \frac{1}{i\not{\partial} + g\not{A}_i + \not{\phi}\gamma_5 + i\hat{m}}, \quad (4)$$

where $\hat{m}$ is the current quark mass matrix: $\text{diag}(m_u, m_d, m_s)$. We assume that the total instanton field A may be approximated as a sum of the single instanton fields, $A = \sum_{i=1}^N A_i$, which is justified with the average size of instantons $\rho \approx 1/3$ fm and average inter-instanton distance $R \approx 1$ fm [28, 29]. Defining the quark propagator $\tilde{S}_0$ with external fields a_μ and the free one S_0 as follows:

$$\tilde{S}_0 = \frac{1}{i\not{\partial} + \not{\phi}\gamma_5 + i\hat{m}}, \quad S_0 = \frac{1}{i\not{\partial} + i\hat{m}}, \quad (5)$$

the $\tilde{S}$ can be expanded with respect to a single instanton:

$$\tilde{S} = \tilde{S}_0 + \sum_i (\tilde{S}_i - \tilde{S}_0) + \sum_{i \neq j} (\tilde{S}_i - \tilde{S}_0) \tilde{S}_0^{-1} (\tilde{S}_j - \tilde{S}_0) + \dots \quad (6)$$

In order to specify the gauge dependence, we rewrite $\tilde{S}_i$ and S_0 in the following form:

$$\begin{aligned} \tilde{S}_i &= L_i S'_i L_i^{-1}, \quad S'_i = \frac{1}{i\not{\partial} + g\not{A}_i + \not{\phi}'\gamma_5 + i\hat{m}}, \\ \tilde{S}_0 &= L_i S'_0 L_i^{-1}, \quad S'_{0i} = \frac{1}{i\not{\partial} + \not{\phi}'\gamma_5 + i\hat{m}}, \end{aligned} \quad (7)$$

where

$$\not{d}'\gamma_5 = L^{-1}(i\not{d} + \not{d}\gamma_5)L. \quad (8)$$

The L denotes the Wilson gauge connection which is expressed as the path-ordered exponent:

$$L_i(x, z_i) = \text{P exp} \left(i \int_{z_i}^x d\xi_\mu a_\mu(\xi) \gamma_5 \right), \quad (9)$$

where z_i denotes an instanton position. Having carried out the manipulation as in Refs. [25, 26], we arrive at the low-frequency part of the fermionic determinant:

$$\begin{aligned} \tilde{\text{Det}}_{\text{low}} &= (\det(i\not{d} + \not{d}\gamma_5 + i\hat{m}))^{-1} \int \prod_f D\psi_f D\psi_f^\dagger \\ &\times e^{\left(\int d^4x \psi_f^\dagger (i\not{d} + \not{d}\gamma_5 + im_f) \psi_f \right)} \prod_f \left\{ \prod_{+}^{N_+} V_{+,f}[\psi_f^\dagger, \psi_f] \prod_{-}^{N_-} V_{-,f}[\psi_f^\dagger, \psi_f] \right\}, \end{aligned} \quad (10)$$

where

$$\tilde{V}_{\pm,f}[\psi_f^\dagger, \psi_f] = \int d^4x \left(\psi_f^\dagger(x) L(x, z) i\not{d}\Phi_{\pm,0}(x; \xi_\pm) \right) \int d^4y \left(\Phi_{\pm,0}^\dagger(y; \xi_\pm) (i\not{d}L(y, z) \psi_f(y) \right) \quad (11)$$

with the fermionic zero-mode solutions $\Phi_{\pm,0}$.

The gauge connection L has some arbitrariness due to its path dependence. However, we can show that such dependence can be minimized. To be more specific, we consider the extended zero mode:

$$(i\not{d} + \not{A} + \not{d}\gamma_5)|\tilde{\Phi}_0\rangle = 0, \quad |\tilde{\Phi}_0^{(1)}\rangle \simeq |\Phi_0\rangle - S_{NZ}\not{d}\gamma_5|\Phi_0\rangle, \quad (12)$$

where A denotes an instanton field located at z , $|\tilde{\Phi}_0^{(1)}\rangle$ stands for the solution in the presence of the external field, and S_{NZ} the well-known non-zero mode of the propagator in the instanton field (see the review [32] and references therein). Here, we have assumed tacitly that the momentum of the external field is small. Inserting the gauge connection $L(x, z)$ into Eq. (12), we obtain:

$$(i\not{d} + \not{A} + \not{d}\gamma_5)|\Phi_0'\rangle = 0, \quad a'_\mu\gamma_5 = L^{-1}(i\partial_\mu + a_\mu\gamma_5)L, \quad |\Phi_0'\rangle = L^{-1}(x, z)|\tilde{\Phi}_0\rangle. \quad (13)$$

If we utilize the relations $(i\not{d} + \not{A})|\Phi_0\rangle = 0$ and $S_{NZ}(i\not{d} + \not{A}) = 1 - |\Phi_0\rangle\langle\Phi_0|$, the solution $|\Phi_0^{(1)'}\rangle$ in Eq. (13) can be reduced to the corresponding solution $|\tilde{\Phi}_0^{(1)}\rangle$ without any problem of the path dependence arising from the gauge connection L .

However, if the zero-mode approximation $S_{NZ} \approx \frac{1}{i\not{d}} = S_{00}$ is used to find the solution ¹

$$|\Phi_{00}^{(1)'}\rangle = |\Phi_0\rangle - S_{00}\not{d}'\gamma_5|\Phi_0\rangle, \quad (14)$$

then

$$|\tilde{\Phi}_{00}^{(1)}\rangle = |\Phi_0\rangle - S_{00}\not{d}\gamma_5|\Phi_0\rangle + S_{00} \left(i \int d\xi_\mu a_\mu(\xi) \gamma_5 \right) |\Phi_0\rangle \quad (15)$$

¹ In fact, the zero-mode approximation is the origin of the problem of the current non-conservation from the instanton vacuum. Pobylytsa suggested a more reliable approximation to restore the current conservation [24].

with applying the inverse gauge connection L^{-1} to Eq. (14). We can easily see that Eq. (15) depends on the path of the gauge connection. We need to choose the path in such a way that $\|\tilde{\Phi}_0^{(1)} - \tilde{\Phi}_{00}^{(1)}\|^2$ can be minimized. As shown explicitly in Ref. [26], the straight line provides the most optimized path.

Having treated the gauge problem in the presence of the external axial-vector field, we finally obtain the low-energy partition function of the nonlocal χ QM from the instanton vacuum:

$$\begin{aligned} \mathcal{Z}[a, m] = & \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{D}\mathcal{M} \exp \int d^4x \left[\psi_f^\dagger(x) (i\not{\partial} + \not{a}\gamma_5 + im_f) \psi_f(x) \right. \\ & \left. + i \int \frac{d^4k d^4p}{(2\pi)^8} e^{-i(k-p)\cdot x} \psi_f^\dagger(k) \sqrt{M_f(k_\mu + a_\mu \gamma_5)} U_{fg}^{\gamma_5} \sqrt{M_g(p_\mu + a_\mu \gamma_5)} \psi_g(p) \right] \end{aligned} \quad (16)$$

with the effective chiral action in the presence of the external axial-vector source

$$\mathcal{S}_{\text{eff}}[a, m] = -N_c \text{Tr} \ln \left[i\not{\partial} + \not{a}\gamma_5 + im_f + i \sqrt{M_f(i\not{\partial}_\mu + a_\mu \gamma_5)} U_{fg}^{\gamma_5} \sqrt{M_g(i\not{\partial}_\mu + a_\mu \gamma_5)} \right], \quad (17)$$

where Tr denotes the functional trace and traces over flavor and spin spaces, generically.

III. MESON DISTRIBUTION AMPLITUDES

The leading-twist pseudoscalar meson light-cone distribution amplitudes (DAs) can be defined with the conserved axial-vector current as follows (here, we consider the light-cone gauge $A \cdot n = 0$):

$$\langle 0 | \bar{q}_f(\tau \hat{n}) \gamma_\mu \gamma_5 q_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle = i\sqrt{2} F_{\mathcal{M}} P_\mu \int_0^1 du e^{i(2u-1)P \cdot \tau \hat{n}} \phi_{\mathcal{M}}(u), \quad (18)$$

where $\mathcal{M}$ denotes the flavor SU(3) octet pseudoscalar meson field with the on-mass shell momentum $P^2 = m_{\mathcal{M}}^2$ in the light-cone frame. In the present work, we choose $\mathcal{M} = \pi^+$ or $\mathcal{M} = K^+$. u and $\hat{n}$ stand for the longitudinal momentum fraction and light-like vector, respectively. $F_{\mathcal{M}}$ is the pseudoscalar meson decay constant. Note that, when the τ approaches to zero, Eq. (18) becomes the pseudoscalar meson decay amplitude due to the DA normalization condition:

$$\int_0^1 du \phi_{\mathcal{M}}(u) = 1. \quad (19)$$

This equation will be used for determining the scale parameter of the model. At this point, we want emphasize that in order to calculate the light-cone DA, it is necessary for us to work in Minkowski space in spite of using the instanton framework which is well-defined in Euclidean space as discussed before. For this purpose, we assume simple analytic continuation (the Wick-rotation) between the two spaces in a practical point of view [12, 13, 15, 27].

The expression for the leading-twist pion and kaon DAs in the present approach is obtained as follows:

$$\phi_{\mathcal{M}}(u) = i \frac{N_c \hat{n}^\mu}{2F_{\mathcal{M}}^2} \int \frac{d^4k}{(2\pi)^4} \delta[uP \cdot \hat{n} - k \cdot \hat{n}] \text{tr}_\gamma \left[\frac{\sqrt{M_f(k)}}{D_f(k)} \gamma_\mu \gamma_5 \frac{\sqrt{M_g(k-P)}}{D_f(k-P)} \gamma_5 \right]$$

$$+ \frac{\sqrt{M_f(k)_\mu} \sqrt{M_g(k-P)}}{D_f(k)} - \frac{\sqrt{M_f(k)} \sqrt{M_g(k-P)_\mu}}{D_f(k-P)} \Big], \quad (20)$$

where $\sqrt{M_f(k)_\mu} = \partial \sqrt{M_f(k)} / \partial k_\mu$. tr_γ denotes the trace over the Dirac spin space. Note that the last two terms that we call the nonlocal contributions in the bracket make the present result gauge-invariant. The quark propagator $D_f(k)$ is expressed in terms of the momentum-dependent dynamical quark mass $M_f(k)$ and current quark mass m_f as follows [31]:

$$D_f(k) = \not{k} - [m_f + M_f(k)]. \quad (21)$$

We employ a simple-pole type parameterization for $F(k)$ in Eq. (3):

$$F(k) = \left[\frac{n\Lambda^2}{(n\Lambda^2 - k^2 + i\epsilon)} \right]^n, \quad (22)$$

since it is easy for the analytic continuation to Minkowski space.

The derivative of the dynamical quark mass $\sqrt{M_\mu}$ appearing in the nonlocal contributions can be evaluated analytically with the simple-pole type form factor of Eq. (22):

$$\sqrt{M_f(k)_\mu} = \frac{1}{2\sqrt{M_f(k)}} \frac{\partial M_f(k)}{\partial k_\mu} = \frac{2n\sqrt{M_f(k)}}{(n\Lambda^2 - k^2)} k_\mu. \quad (23)$$

Inserting Eqs. (22) and (23) into Eq. (20), we finally obtain the expression for the leading-twist light-cone pion and kaon DAs:

$$\begin{aligned} \phi_{\mathcal{M}}(u) &= i \frac{N_c}{2F_{\mathcal{M}}^2} \int \frac{d^4k}{(2\pi)^4} \delta[uP_+ - k_+] \sqrt{M_f(k)} \sqrt{M_f(k-P)} \text{tr}_\gamma \left[\frac{1}{D_f(k)} \not{k} \gamma_5 \frac{1}{D_g(k-P)} \gamma_5 \right. \\ &\quad \left. + \frac{2nk_+}{D_f(k)(n\Lambda^2 - k^2)} - \frac{2n(k_+ - P_+)}{D_g(k-P)[n\Lambda^2 - (k-P)^2]} \right] \\ &= \frac{N_c}{2F_{\mathcal{M}}^2} \int \frac{dk_- dk_T^2}{(2\pi)^3} [\mathcal{T}_L + \mathcal{T}_{NL}], \end{aligned} \quad (24)$$

where p_+ stands for $p \cdot \hat{n}$. The subscripts L and NL stand for the local and nonlocal contributions, respectively. Further evaluation of $\mathcal{T}_{L,NL}$ can be found in Appendix.

IV. RESULTS AND DISCUSSION

In this Section, we discuss the numerical results for the leading-twist light-cone pion and kaon DAs with the gauged nonlocal effective chiral action. We first fix the scale parameter of the model, i.e. Λ in Eq. (22). In Ref. [14], the Λ was chosen to be 1.2 GeV by using the normalization condition in Eq. (19), with which the pion and kaon decay constants are simultaneously reproduced. However, since the Ward-Takahashi identity for the axial-vector current is broken, the improper Pagels-Stokar (PS) expression for the pion decay constant had to be used to normalize the DAs properly. In the present gauge-invariant approach, we are able to determine the scale parameter Λ without any trouble.

In Table I, we list the pion and kaon decay constants evaluated by using the normalization condition in Eq. (19). As shown in Table I, when we set $\Lambda \simeq 1.0$ GeV, the values of

F_π are in accord with the empirical one ($F_\pi = 93$ MeV) for powers of the form factor $n = 1, 2, 3$. We find that without the nonlocal contribution from gauge invariance, the results are underestimated by about $10 \sim 20\%$. The kaon decay constant F_K is found to be about 10% smaller than the empirical data $F_K = 113$ MeV. However, it is well known that the meson-loop corrections play a crucial role in describing the F_K [33].

	F_π (Exp: 93 MeV)			F_K (Exp: 113 MeV)		
n	Local	Nonlocal	Total	Local	Nonlocal	Total
1	87.03	28.75	91.66	92.59	31.94	97.95
2	84.47	30.38	89.77	90.87	34.37	97.15
3	83.68	31.02	89.24	90.31	35.31	96.97

TABLE I: The results of the pion and kaon decay constants F_π and F_K [MeV].

In Fig. 1, we depict the pion and kaon DAs in the upper and lower panels, respectively. The dashed, long dashed, and solid curves represent the local, nonlocal, and total contributions to the DAs, respectively. For comparison, we also draw the asymptotic DA in the dotted curve in each panel. As the power of the form factor n increases, the local contributions are almost independent of n , whereas the nonlocal ones are drastically changed: Their humped shape is getting manifest as n increases. As a result, the total pion DA is getting flattened in the region of $0.25 \lesssim u \lesssim 0.75$, as n increases. Although we do not present explicitly the pion DA with higher n , it is even more flattened, if we increase n higher than three. Note that the pion DA is suppressed at the end points of u . In the lower panels of Fig. 1, we find that while the local contribution to the kaon DA is almost symmetric, the nonlocal one turns out to be rather asymmetric. However, when these two contributions are added, the kaon DA is unexpectedly similar to the asymptotic one. However, the kaon DA is quite much suppressed at the end points, in particular, with higher n , compared to the asymptotic one. It indicates that the effects of the current quark mass becomes obvious for the kaon DA at the end points, as shown in Ref. [14]. The suppression of the pion and kaon DAs at the end points corresponds to the negative second Gegenbauer moments.

It is also of great use to examine the Gegenbauer polynomial to analyze the DAs in detail [14, 15]. The Gegenbauer moments for the DAs are defined by the coefficients of the Gegenbauer polynomial expansion of the DAs, a_m :

$$\phi(u) = 6u(1-u)[1 + a_1 C_1^{3/2}(\xi) + a_2 C_2^{3/2}(\xi) + \cdots]. \quad (25)$$

Note that the odd $(2m+1)$ Gegenbauer moments are all zero for the pion DA because of isospin symmetry. The Gegenbauer moments to the sixth order ($m=6$) are evaluated and are listed in Table II for the pion and in Table III for the kaon.

Our results of a_2^π for $n=1$ are rather consistent with those of various model calculations listed in Table II. We find that the absolute values of a_2^π are in general larger than those given in Ref. [14] in which the nonlocal contributions were not considered. Moreover, we verified that the difference between those with and without the nonlocal ones becomes evident as n increases. We obtain the negative values of a_4^π for $n=2$ and 3, which are different from those of other models [17, 37]. As mentioned previously, the sign for a_4 plays an important role in the end-point behavior of the pion and kaon DAs. The negative sign indicates that the DAs are suppressed at the end points, i.e. the DAs turn out to be in concave shapes at the end points.




FIG. 1: Pion (upper panels) and kaon (lower panels) distribution amplitudes. The dashed, long dashed, and solid curves represent the local, nonlocal, and total contributions, respectively. The asymptotic DA is also drawn for comparison.

In Table III, we list the Gegenbauer moments for the kaon DA, including both odd and even ones. The most interesting feature shown here is that the signs of the calculated a_1^K are all negative. The results of the QCD sum rule calculations are not conclusive in determining the sign of a_1^K [34, 36, 38, 39]. An interesting discussion on this issue can be found in Ref. [34]. In addition, the a_2^K are also negative for $n = 2, 3$, being obviously different from the pion. Note that the values of a_2^K are all positive for the QCDSR calculations in Refs. [35, 36, 37].

n	a_2^π	a_4^π	a_6^π
1	0.11911	0.01438	-0.00009
2	0.07127	-0.03605	-0.02103
3	0.05339	-0.06088	-0.02596
[4]	0.56	—	—
[17] (2.4 GeV)	—	0.044 ± 0.016	0.023 ± 0.010
[19] (2.4 GeV)	0.12 ± 0.03	—	—
[34] (1.0 GeV)	$0.26^{+0.21}_{-0.09}$	—	—
[35] (2.24 GeV)	0.236(82)	—	—
[36] (1.0 GeV)	0.25 ± 0.15	—	—
[37] (1.0 GeV)	0.44	0.25	—

TABLE II: The Gegenbauer moments for the pion DA. The numbers in the parentheses stand for the renormalization scale.

The analysis of the Gegenbauer moments a_2^π and a_4^π has been performed first by Schmedding and Yakovlev [19], based on the CLEO data of the transition form factor $F_{\gamma^*\gamma\pi}(Q^2)$ at the renormalization point $\mu = 2.4$ GeV (SY-scale). In order to compare them to the present results of the Gegenbauer moments, it is necessary to evolve the scale of the present calculation to the SY-scale by using the one-loop QCD evolution equation. Note that the anomalous dimension for the Gegenbauer moments for the leading-twist DAs are obtained as follows [1, 2, 3, 4]:

$$\gamma_m^{(0)} = -\frac{8}{3} \left[3 + \frac{2}{(m+1)(m+2)} - 4 \sum_{m'=1}^{m+1} \frac{1}{m'} \right], \quad (26)$$

where $\gamma_1^{(0)} = 7.11$, $\gamma_2^{(0)} = 11.11$, $\gamma_3^{(0)} = 13.95$, $\gamma_4^{(0)} = 16.18$. Thus, the Gegenbauer moments in two different renormalization scales, $\Lambda_1 = 2.4$ GeV and $\Lambda_2 = 1.0$ GeV, can be compared by the following relation:

$$a_m(\Lambda_1) = a_m(\Lambda_2) \left[\frac{\alpha(\Lambda_1)}{\alpha(\Lambda_2)} \right]^{\gamma_m^{(0)}/(2\beta_0)} \sim a_m(\Lambda_2) \left[\frac{\ln[\Lambda_2/\Lambda_{\text{QCD}}]}{\ln[\Lambda_1/\Lambda_{\text{QCD}}]} \right]^{\gamma_m^{(0)}/(2\beta_0)}. \quad (27)$$

Here, we use $\beta_0 = 9$ and $\Lambda_{\text{QCD}} \simeq 0.2$ GeV. The contours on the Gegenbauer parameter plane (a_2, a_4) analyzed from the CLEO experiment are depicted together with our results in Fig. 2. The solid and dashed ellipses indicate the 68% and 95% confidential levels, respectively [19]. We also indicate the positions for the asymptotic DA (AS) and for the Chenyak-Zhitnitsky (CZ) DA [40]. The closed circles ($\bullet$) denote the values of (a_2^π, a_4^π) calculated in the present work. On the contrary, the open circles ($\circ$) are taken from Ref. [14] in which the gauge invariance of the effective chiral action was not considered. We observe that the closed circles for $n = 1, 2, 3$ are all inside the 95% confidential levels. However, when the nonlocal contributions are turned off (open circles), which is identical to the results of Ref. [14], a_2^π and a_4^π for $n = 3$ lie outside the 95% confidential regions. We also see that the values of $|a_4^\pi|$ become slightly larger with the nonlocal contributions. We infer from it that the pion DA is getting more flattened by larger values of $|a_4^\pi|$.

In addition, we depict the results for $\Lambda = 1.2$ GeV and 1.5 GeV in the stars ($\star$) and asterisks ($*$), respectively in Fig. 2, to test the dependence on the scale parameter of the model Λ . We see that as Λ increases the positions of (a_2^π, a_4^π) move toward the center of the 95% confidential level. The a_4^π is less sensitive to the Λ , compared to a_2^π . We verified that

n	a_1^K	a_2^K	a_3^K	a_4^K	a_5^K	a_6^K
1	-0.01879	0.02525	0.00467	0.00676	-0.00283	0.00069
2	-0.01193	-0.01935	0.00599	-0.02470	-0.00371	-0.01183
3	-0.01012	-0.03694	0.00706	-0.03840	-0.00566	-0.01365
[34] (1.0 GeV)	$0.07^{+0.02}_{-0.03}$	$0.27^{+0.37}_{-0.12}$	—	—	—	—
[36] (1.0 GeV)		0.30 ± 0.15	—	—	—	—
[38] (1.0 GeV)	-0.18 ± 0.09	0.16 ± 1.10	—	—	—	—
[39] (1.0 GeV)	0.10 ± 0.12	—	—	—	—	—

TABLE III: The Gegenbauer moments for the kaon DA. The numbers in the parentheses stand for the renormalization scale.




FIG. 2: Gegenbauer parameter space ($a_2 - a_4$) for the pion DA with the CLEO experimental data analysis taken from Ref. [18, 19] in the renormalization scale $\mu = 2.4\text{ GeV}$. The solid and dashed ellipses indicate the 68% (1σ) and 95% (2σ) confidential levels of the analysis. The open circles (o) denote the results of the Gegenbauer moments without the nonlocal contributions ($\Lambda = 1.2\text{ GeV}$) [14]. The close circles ($\bullet$), stars ($\star$), and asterisks ($*$) stand for the cases with the nonlocal contributions for $\Lambda = 1.0\text{ GeV}$, 1.2 GeV and 1.5 GeV , respectively. The results with the same power n and different Λ are connected by the solid line.

the calculated pion decay constants F_π for $\Lambda = 1.2\text{ GeV}$ and 1.5 GeV are only smaller by about 10% than the empirical one.

To show the dependence of the pion DAs on the scale parameter Λ , we draw in Fig. 3 the pion DA with different Λ . We observe that the pion DAs are more suppressed at the end points and more humped as Λ increases.




FIG. 3: Pion distribution amplitudes at $\mu = 2.4\text{ GeV}$ with $\Lambda = 1.0\text{ GeV}$, 1.2 GeV and 1.5 GeV from the left end panel.

V. SUMMARY AND CONCLUSION

We investigated the pion and kaon distribution amplitudes within the framework of the nonlocal chiral quark model from the instanton vacuum, making the gauge invariance of the low-energy partition function preserved in the presence of the external fields. The scale parameter of the present work was set to be $\Lambda \simeq 1.0$ GeV which is not far from that determined by using the saddle-point equation. Using $\Lambda \simeq 1.0$ GeV, we were able to reproduce the empirical values of the pion and kaon decay constants qualitatively well by using the normalization condition of the distribution amplitudes. The effects of flavor SU(3) symmetry breaking effects were explicitly taken into account consistently.

We found that the nonlocal contribution arising from the gauge invariance of the effective chiral action makes the pion distribution amplitude flattened more in the region of $0.25 \lesssim u \lesssim 0.75$, compared to the former investigation without consideration of the gauge invariance [14]. The results indicate that the absolute values of Gegenbauer moments a_2 and a_4 become larger. As for the kaon, the nonlocal contribution is less important. We noticed that a_1^K is still negative even with the nonlocal contributions, which is similar to those in the previous work [14].

By virtue of the renormalization group equation, we evolved the Gegenbauer moments a_2 and a_4 so that we may compare them with the Schmedding-Yakovlev analysis on the CLEO experimental data at $\mu = 2.4$ GeV. We found that the present results with the gauge invariance turn out to satisfy the Schmedding-Yakovlev analysis of the Gegenbauer moments at the 95% confidential level. We found that the values of (a_2^K, a_4^K) were very close to those of the asymptotic distribution amplitude.

In conclusion, the present gauge-invariant approach has several advantages, compared to the former works. Firstly, it provides a more reasonable scale to the pion and kaon DAs, since we can use the normalization condition without any gauge-invariant problem. Secondly, the nonlocal corrections arising from the gauge invariance of the effective chiral action play a significant role in describing the pion distribution amplitude. Compared to the Schmedding-Yakovlev analysis, the present results are phenomenologically better than those of the former work. In order to describe the pion and kaon distribution amplitudes more consistently, we want to introduce the meson-loop corrections ($1/N_c$ corrections) to them. The corresponding work is under progress.

Acknowledgments

The work of S.N. is supported by the Brain Korea 21 (BK21) project in Center of Excellency for Developing Physics Researchers of Pusan National University, Korea. S.N. would like to thank J. H. Lee for fruitful discussions.

Appendix

The first term in the square parenthesis in Eq. (24) reads

$$\text{tr}_\gamma \left[\frac{\sqrt{M_f(k)}}{D(k)} \not{p} \gamma_5 \frac{\sqrt{M_f(k-P)}}{D(k-P)} \gamma_5 \right] = -4M_0 \sqrt{f(m_f)f(m_f)} \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^n \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^n$$

$$\begin{aligned}
& \times \left\{ k \cdot \hat{n} \left[m_g + M_0 f(m_g) \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{2n} \right] - (k-P) \cdot \hat{n} \left[m_f + M_0 f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} \right] \right\} \\
& \times \left\{ k^2 - m_f^2 - 2M_0 m_f f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} - M_0^2 f^2 \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{4n} \right\}^{-1} \\
& \times \left\{ (k-P)^2 - m_g^2 - 2M_0 m_g f(m_g) \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{2n} - M_0^2 f^2 \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{4n} \right\}^{-1} \\
& = 4\sqrt{\eta_f \eta_g} (\alpha k_- - \gamma_f)^n (\beta k_- - \gamma_g)^n \\
& \times \frac{\alpha (\alpha k_- - \gamma_f)^{2n} [m_g (\beta k_- - \gamma_g)^{2n} + \eta_g] - \beta (\beta k_- - \gamma_g)^{2n} [m_f (\alpha k_- - \gamma_f)^{2n} + \eta_f]}{\mathcal{D}_f \mathcal{D}_g},
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{D}_f &= (\alpha k_- - D_f)(\alpha k_- - \gamma_f)^{4n} - 2\eta_f m_f (\alpha k_- - \gamma_f)^{2n} - \eta_f^2 + i\epsilon, \\
\mathcal{D}_g &= (\beta k_- - D_g)(\beta k_- - \gamma_g)^{4n} - 2\eta_g m_g (\beta k_- - \gamma_g)^{2n} - \eta_g^2 + i\epsilon.
\end{aligned}$$

We use the following parameterizations for convenience:

$$\begin{aligned}
k^2 &= k_+ k_- - k_T^2 = u P_+ k_- - k_T^2 = \alpha k_- - k_T^2, \\
(k-P)^2 &= (u-1) P_+ k_- - (u-1) m_\phi^2 - k_T^2 = \beta k_- - (u-1) m_\phi^2 - k_T^2, \\
n\Lambda^2 - k^2 &= -u P_+ k_- + [k_T^2 + n\Lambda^2] = -\alpha k_- + \gamma_f, \\
n\Lambda^2 - (k-P)^2 &= -(u-1) P_+ k_- + [(u-1) m_\phi^2 + k_T^2 + n\Lambda^2] = -\beta k_- + \gamma_g, \\
k^2 - m_f^2 &= \alpha k_- - [k_T^2 + m_f^2] = \alpha k_- - \delta_f, \\
(k-P)^2 - m_g^2 &= \beta k_- - [(u-1) m_\phi^2 + k_T^2 + m_g^2] = \beta k_- - \delta_g, \\
M_0 f(m_f) (n\Lambda^2)^{2n} &= \eta_f, \quad M_0 f(m_g) (n\Lambda^2)^{2n} = \eta_g, \\
M_f(k) &= \frac{\eta_f}{(\alpha k_- - \gamma_f)^{2n}}, \quad M_f(k-P) = \frac{\eta_g}{(\beta k_- - \gamma_g)^{2n}}, \\
\frac{1}{D(k)} &= \frac{(\alpha k_- - \gamma_f)^{4n}}{\mathcal{D}_f} \left[k + m_f + \frac{\eta_f}{(\alpha k_- - \gamma_f)^{2n}} \right], \\
\frac{1}{D(k-P)} &= \frac{(\beta k_- - \gamma_g)^{4n}}{\mathcal{D}_g} \left[(k-P) + m_g + \frac{\eta_g}{(\beta k_- - \gamma_g)^{2n}} \right].
\end{aligned}$$

The second term can be evaluated using the dynamical quark mass of Eq. (22) as follows:

$$\begin{aligned}
& 2n \text{tr}_\gamma \left[\frac{\hat{n} \cdot k \sqrt{M_f(k)} \sqrt{M_f(k-P)}}{D(k)(n\Lambda^2 - k^2)} \right] = \frac{8nk \cdot \hat{n} M_0 \sqrt{f(m_f) f(m_g)}}{(n\Lambda^2 - k^2)} \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^n \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^n \\
& \times \left\{ m_f + M_0 f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} \right\} \left\{ k^2 - m_f^2 - 2M_0 f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} - M_0^2 f^2 \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{4n} \right\}^{-1} \\
& = -\frac{8n\alpha \sqrt{\eta_f \eta_g} [m_f (\alpha k_- - \gamma_f)^{2n} + E_f] (\alpha k_- - \gamma_f)^{n-1}}{(\beta k_- - \gamma_g)^n \mathcal{D}_f},
\end{aligned}$$

Similarly, the third term can be written as follows:

$$2n\text{tr}_\gamma \left[\frac{(k-P) \cdot \hat{n} \sqrt{M_f(k)} \sqrt{M_f(k-P)}}{D(k-P)[n\Lambda^2 - (k-P)^2]} \right] = - \frac{8n\beta \sqrt{\eta_f \eta_g} [m_g(\beta k_- - \gamma_g)^{2n} + \eta_g] (\beta k_- - \gamma_g)^{n-1}}{(\alpha k_- - \gamma_f)^n \mathcal{D}_g}.$$

For convenience, we denote these three terms evaluated above by $\mathcal{T}_L^V$ and $\mathcal{T}_{NL}^V$, which is the sum of the second and third terms.

-
- [1] A. V. Efremov and A. V. Radyushkin, Phys. Lett. B **94**, 245 (1980).
 - [2] G. P. Lepage and S. J. Brodsky, Phys. Lett. B **87**, 359 (1979).
 - [3] G. P. Lepage and S. J. Brodsky, Phys. Rev. D **22**, 2157 (1980).
 - [4] V. L. Chernyak and A. R. Zhitnitsky, Phys. Rept. **112**, 173 (1984).
 - [5] V. M. Braun and I. E. Filyanov, Z. Phys. C **44**, 157 (1989).
 - [6] A. P. Bakulev and S. V. Mikhailov, Z. Phys. C **68**, 451 (1995).
 - [7] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Lett. B **508**, 279 (2001) [Erratum-ibid. B **590**, 309 (2004)].
 - [8] A. P. Bakulev and N. G. Stefanis, Nucl. Phys. B **721**, 50 (2005).
 - [9] P. Ball and A. N. Talbot, JHEP **0506**, 063 (2005).
 - [10] L. Del Debbio, M. Di Pierro and A. Dougall, Nucl. Phys. Proc. Suppl. **119**, 416 (2003).
 - [11] S. Dalley and B. van de Sande, Phys. Rev. D **67**, 114507 (2003).
 - [12] V. Y. Petrov, M. V. Polyakov, R. Ruskov, C. Weiss and K. Goeke, Phys. Rev. D **59**, 114018 (1999).
 - [13] A. E. Dorokhov, JETP Lett. **77**, 63 (2003) [Pisma Zh. Eksp. Teor. Fiz. **77**, 68 (2003)].
 - [14] S. i. Nam, H. C. Kim, A. Hosaka and M. M. Musakhanov, Phys. Rev. D **74**, 014019 (2006).
 - [15] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **64**, 074003 (2001).
 - [16] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **66**, 054002 (2002).
 - [17] E. Ruiz Arriola and W. Broniowski, Phys. Rev. D **66**, 094016 (2002).
 - [18] J. Gronberg *et al.* [CLEO Collaboration], Phys. Rev. D **57**, 33 (1998).
 - [19] A. Schmedding and O. I. Yakovlev, Phys. Rev. D **62**, 116002 (2000).
 - [20] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Rev. D **67**, 074012 (2003).
 - [21] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Lett. B **578**, 91 (2004).
 - [22] M. Chretien and R. E. Peierls, Proc. Roy. Soc. Lond. A **223**, 468 (1954).
 - [23] B. Holdom, J. Terning, and K. Verbeek, Phys. Lett. B **232**, 351 (1989).
 - [24] P. V. Pobylitsa, Phys. Lett. B **226**, 387 (1989).
 - [25] M. M. Musakhanov and H.-Ch. Kim, Phys. Lett. B **572**, 181 (2003).
 - [26] H.-Ch. Kim, M. Musakhanov and M. Siddikov, Phys. Lett. B **608**, 95 (2005).
 - [27] I. V. Anikin, A. E. Dorokhov and L. Tomio, Phys. Atom. Nucl. **64**, 1329 (2001) [Yad. Fiz. **64**, 1405 (2001)].
 - [28] E. V. Shuryak, Nucl. Phys. B **203**, 93 (1982).
 - [29] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **272**, 457 (1986).
 - [30] M. Musakhanov, Eur. Phys. J. C **9**, 235 (1999).
 - [31] M. Musakhanov, Nucl. Phys. A **699**, 340 (2002).
 - [32] T. Schafer and E. V. Shuryak, Rev. Mod. Phys. **70**, 323 (1998).
 - [33] J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985).

- [34] A. Khodjamirian, T. Mannel and M. Melcher, Phys. Rev. D **70**, 094002 (2004).
- [35] M. Gockeler *et al.*, hep-lat/0510089.
- [36] P. Ball, V. M. Braun and A. Lenz, JHEP **0605**, 004 (2006).
- [37] V. M. Braun and I. E. Filyanov, Z. Phys. C **48**, 239 (1990) [Sov. J. Nucl. Phys. **52**, 126 (1990 YAFIA,52,199-213.1990)].
- [38] P. Ball and M. Boglione, Phys. Rev. D **68**, 094006 (2003).
- [39] V. M. Braun and A. Lenz, Phys. Rev. D **70**, 074020 (2004).
- [40] V. L. Chernyak and A. R. Zhitnitsky, JETP Lett. **25**, 510 (1977) [Pisma Zh. Eksp. Teor. Fiz. **25**, 544 (1977)].